

Available online at www.sciencedirect.com

Resuscitation

journal homepage: www.elsevier.com/locate/resuscitation

Clinical paper

Association between time of day and CPR quality as measured by CPR hemodynamics during pediatric in-hospital CPR



EUROPEAN

RESUSCITATION

Heather A. Wolfe^{a,*}, Ryan W. Morgan^a, Robert M. Sutton^a, Ron W. Reeder^b, Kathleen L. Meert^c, Murray M. Pollack^{d,k}, Andrew R. Yates^e, John T. Berger^d, Christopher J. Newth^f, Joseph A. Carcillo^g, Patrick S. McQuillen^h, Rick E. Harrisonⁱ, Frank W. Moler^j, Todd C. Carpenter^l, Daniel A Notterman^m, J. Michael Dean^b, Vinay M. Nadkarni^a, Robert A. Berg^a, for the Eunice Kennedy Shriver National Institute of Child Health Human Development Collaborative Pediatric Critical Care Research Network Pediatric Intensive Care Quality of Cardiopulmonary Resuscitation (PICqCPR) investigators

^a Department of Anesthesiology and Critical Care Medicine, The Children's Hospital of Philadelphia, University of Pennsylvania, Philadelphia, PA, United States

- ^b Department of Pediatrics, University of Utah, Salt Lake City, Utah, United States
- ^c Department of Pediatrics, Children's Hospital of Michigan, Wayne State University, Detroit, MI, United States

^d Department of Pediatrics, Children's National Medical Center, Washington D.C., United States

e Department of Pediatrics, Nationwide Children's Hospital, The Ohio State University, Columbus, OH, United States

[†] Department of Anesthesiology, Children's Hospital Los Angeles, University of Southern California Keck School of Medicine, Los Angeles, CA, United States

- ⁹ Department of Critical Care Medicine, Children's Hospital of Pittsburgh, University of Pittsburgh, Pittsburgh, PA, United States
- ^h Department of Pediatrics, Benioff Children's Hospital, University of California San Francisco, San Francisco, CA, United States
- ¹ Department of Pediatrics, Mattel Children's Hospital, University of California Los Angeles, Los Angeles, CA, United States

¹ Department of Pediatrics, C.S. Mott Children's Hospital, University of Michigan, MI, United States

- ^k Department of Pediatrics, Phoenix Children's Hospital, Phoenix, AZ, United States
- ¹ Department of Pediatrics, Denver Children's Hospital, University of Colorado, Denver, CO, United States
- ^m Department of Molecular Biology, Princeton University, Princeton, New Jersey, United States

Abstract

Introduction: Patients who suffer in-hospital cardiac arrest (IHCA) are less likely to survive if the arrest occurs during nighttime versus daytime. Diastolic blood pressure (DBP) as a measure of chest compression quality was associated with survival from pediatric IHCA. We hypothesized that DBP during CPR for IHCA is lower during nighttime versus daytime.

Methods: This is a secondary analysis of data collected from the Pediatric Intensive Care Quality of Cardiopulmonary Resuscitation Study. Pediatric or Pediatric Cardiac Intensive Care Unit patients who received chest compressions for \geq 1 min and who had invasive arterial BP monitoring were enrolled. Nighttime was defined as 11:00PM to 6:59AM and daytime as 7:00AM until 10:59PM. Primary outcome was attainment of DBP \geq 25 mmHg in infants

E-mail address: wolfeh@email.chop.edu (H.A. Wolfe).

https://doi.org/10.1016/j.resuscitation.2020.06.027

Received 2 March 2020; Received in revised form 4 June 2020; Accepted 16 June 2020 0300-9572/© 2020 Elsevier B.V. All rights reserved.

^{*} Corresponding author at: The University of Pennsylvania Perelman School of Medicine, The Children's Hospital of Philadelphia, 3401 Civic Center Blvd, 6th Floor Wood Building Rm 6040, Philadelphia, PA 19104-4399, United States.

<1 year and \geq 30 mmHg in older children. Secondary outcomes were mean DBP, ROSC, and survival to hospital discharge. Univariable and multivariate analyses evaluated the relationships between time (nighttime vs. daytime) and outcomes.

Results: Between July 1, 2013 and June 30, 2016, 164 arrests met all inclusion/exclusion criteria: 45(27%) occurred at nighttime and 119(73%) during daytime. Average DBPs achieved were not different between groups (DBP: nighttime 28.3 mmHg[25.3, 36.5] vs. daytime 29.6 mmHg[21.8, 38.0], p = 0.64). Relative risk of DBP threshold met during nighttime vs. daytime was 1.27, 95%CI [0.80, 1.98], p = 0.30. There was no significant nighttime vs. daytime difference in ROSC (28/45[62%] vs. 84/119[71%] p = 0.35) or survival to hospital discharge (16/45[36%] vs. 61/119[51%], p = 0.08). **Conclusions:** In this cohort of pediatric ICU patients with IHCA, there was no significant difference in DBP during CPR between nighttime and daytime. **Keywords:** Cardiopulmonary resuscitation (CPR), Cardiac arrest, Pediatric, In-Hospital, Survival, Outcomes

Introduction

More than 15,000 children receive in-hospital CPR each year in the United States,¹ and the ratio of intensive care unit (ICU) to ward events is 19 to 1.² Overall, more than 1% of children admitted to a pediatric intensive care unit have a cardiac arrest in the unit, with the frequency approximately three-fold higher in pediatric cardiac intensive care units.^{3,4} While survival rates in the United States have improved over the past decades, most children who suffer from an IHCA do not survive to hospital discharge.⁵

Large studies have consistently established that survival rates are lower following cardiac arrests at nighttime compared with daytime for both in-hospital and out-of-hospital cardiac arrests in children and adults.^{6–12} Investigators have postulated that the worse outcomes may be due to worse CPR quality or worse post-resuscitation care at night, and that either could result from lower nighttime staffing or nighttime fatigue among providers. However, these studies emanate from large in-hospital and out-of-hospital cardiac arrest databases that lack data on CPR quality.

The Pediatric Intensive Care Quality of CPR (PICqCPR) study was a multi-institution investigation of 164 children with invasive hemodynamic monitoring during CPR. Consistent with multiple translational laboratory studies the PICqCPR data demonstrated that mean diastolic blood pressure (DBP) of > 25 mmHg in infants and > 30 mmHg children during CPR was associated with greater likelihood of survival to discharge and survival with favorable neurologic outcome.13 Because the number of patients in the PICqCPR study was deemed to be insufficient to address daytime versus nighttime survival outcomes, our goal was to evaluate whether CPR quality differed at nighttime compared to daytime as evidenced by DBP attained during CPR, using this rich CPR hemodynamic database. Our primary hypothesis was that CPR quality is worse at nighttime. Our primary outcome measure was attainment of mean DBP \geq 25 mmHg in infants and \geq 30 mmHg in children, and our secondary outcomes were overall mean DBP during CPR, return of spontaneous circulation (ROSC) and survival to discharge.

Methods

The PICqCPR study was a prospective multicenter cohort study of CPR occurring in the ICUs of 11 institutions in the CPCCRN network between July 1, 2013 and June 30, 2016.¹³ This study is a secondary observational analysis of the original multi-center cohort study, which was approved with waiver of informed consent by the Institutional Review Boards at the University of Utah Data Coordinating Center (DCC) and at each participating clinical site. CPCCRN research coordinators obtained Utstein-style standardized cardiac arrest and

CPR data,^{14,15} including demographics, preexisting conditions, illness categories, interventions in place at time of arrest, first documented rhythm, immediate cause of arrest, duration of CPR, resuscitation interventions, and outcome data, with assessments of neurological outcomes by Pediatric Cerebral Performance Categories (PCPC) and functional status by Functional Status Scale (FSS)¹⁶ scores for pre-admission status and at hospital discharge. The FSS is scored over 6 domains with a total score from 6 points up to a maximum of 30 points, with higher values indicating worse outcomes.¹⁶

In order to be included in the parent study, patients were required to have an arterial line in place at the time of arrest. Arterial BP waveforms were printed from study sites, de-identified, and stored at the DCC. Waveforms were analyzed by Children's Hospital of Philadelphia investigators who were blinded to patient outcomes (PlotDigitizer; Version 2.0; Department of Physics, University of South Alabama).

Patient population

Patients in a CPCCRN ICU \geq 37 weeks' gestation and <19 years old who received chest compressions for \geq 1 min and had invasive arterial BP monitoring during CPR were eligible for inclusion. In addition, patients were required to have central venous pressures, respiratory plethysmography or electrocardiographic waveform data sufficient to determine stops and starts in CPR. Patients were excluded if the first chest compression of CPR was not captured, if they did not have at least 1 min of continuous arterial BP waveform, or if CPR starts and stops could not be determined. Research coordinators identified potential cardiac arrest patients via paging systems and daily screening of the ICU census.

Time of day and day of week definitions

Nighttime was defined as 11:00 PM to \leq 6:59 AM and daytime was defined as 7:00 AM until 10:59 PM, as per previous investigations.^{6,8,11} Similarly, weekdays were defined as 7:00 AM Monday until 10:59 PM Friday, weekends were defined as 10:59 PM Friday until 7:00 AM on Monday.

Blood pressure determination

For each minute of CPR, a mean DBP and mean SBP were determined. Subsequently, mean DBP and mean SBP for each patient event was calculated as the average BP over the first 10 min of CPR. If the patient received <10 min of CPR, the mean BPs could only be determined for the minutes of CPR performed. The original publication has further details regarding parent study methods and arterial blood pressure data determination.¹³ The DBP threshold was

defined as mean DBP \geq 25 mmHg during CPR in infants and \geq 30 mmHg in children \geq 1 year old. Systolic BP threshold was defined as mean SBP \geq 60 mmHg during CPR in infants and \geq 80 mmHg in children \geq 1 year old. For the first 20 min post-ROSC, SBP and DBP values were measured at 5-min intervals. Systolic and diastolic hypotension were defined as any one of these 5-min values being below the age-based 5th percentile for SBP and DBP, respectively based on normative data for age, sex and height.¹⁷

Outcomes

Our primary outcome measure was attainment of DBP associated with improved outcomes (mean DBP \geq 25 mmHg in infants and \geq 30 mmHg in older children), and our secondary outcomes were overall mean DBP, ROSC, and survival to discharge. We also assessed intraarrest SBP, post-ROSC hypotension, 24-h survival, survival with favorable neurologic status, and change in FSS from baseline. New

morbidity at hospital discharge is defined as an increase in FSS of at least 3 points or increase of 2 in a single FSS domain. 16

Statistical analysis

Standard descriptive statistics were used to summarize patient and event characteristics. Differences in characteristics and outcomes between patients who received CPR during the nighttime versus daytime as well as weekday versus weekend were examined using Fisher's exact test for categorical variables or the Wilcoxon rank-sum test for continuous variables. Primary and secondary outcomes were further compared between groups using Poisson regression models with robust error estimates, adjusting for age category (<1 year, \geq 1 year), first documented rhythm (asystole/PEA,VF/VT, bradycardia with poor perfusion), illness category (surgical cardiac, medical cardiac, surgical non-cardiac, medical non-cardiac), and location of CPR (pediatric ICU, pediatric cardiac ICU). These results were

Table 1 – Patient Characteristics by CPR Time of Day.

| | Time of CPR | | | | |
|----------------------------|------------------|---------------------------|----------------------------|-------------------|--|
| | Overall(N = 164) | Night ^e (N=45) | Day ^e (N = 119) | p-Value | |
| Age (Years) | 0.0 [0.0, 2.5] | 0 [0,3] | 0 [0,2] | 0.55 ^b | |
| Age Category | | | | 0.93 ^a | |
| <1 month | 41 (25%) | 10 (22%) | 31 (26%) | | |
| 1 month - <1 year | 57 (35%) | 16 (36%) | 41 (34%) | | |
| 1 year - <8 years | 41 (25%) | 11 (24%) | 30 (25%) | | |
| 8 years - <19 years | 25 (15%) | 8 (18%) | 17 (14%) | | |
| Male | 90 (55%) | 25 (56%) | 65 (55%) | 1.00 ^a | |
| Race | | | | 0.18 ^a | |
| White | 82 (50%) | 26 (58%) | 56 (47%) | | |
| Black or African American | 37 (23%) | 12 (27%) | 25 (21%) | | |
| Other | 8 (5%) | 0 (0%) | 8 (7%) | | |
| Not Reported | 37 (23%) | 7 (16%) | 30 (25%) | | |
| Preexisting conditions | | | | | |
| Respiratory insufficiency | 132 (80%) | 38 (84%) | 94 (79%) | 0.51 ^ª | |
| Hypotension | 128 (78%) | 37 (82%) | 91 (76%) | 0.53 ^a | |
| Congestive heart failure | 19 (12%) | 6 (13%) | 13 (11%) | 0.79 ^a | |
| Pneumonia | 13 (8%) | 3 (7%) | 10 (8%) | 1.00 ^a | |
| Sepsis | 44 (27%) | 10 (22%) | 34 (29%) | 0.55 ^a | |
| Renal insufficiency | 24 (15%) | 6 (13%) | 18 (15%) | 1.00 ^a | |
| Malignancy | 5 (3%) | 2 (4%) | 3 (3%) | 0.62 ^a | |
| Congenital heart disease | 99 (60%) | 29 (64%) | 70 (59%) | 0.59 ^a | |
| Illness Category | | | | 0.76 ^a | |
| Surgical cardiac | 88 (54%) | 24 (53%) | 64 (54%) | | |
| Medical cardiac | 25 (15%) | 8 (18%) | 17 (14%) | | |
| Surgical non-cardiac | 13 (8%) | 2 (4%) | 11 (9%) | | |
| Medical non-cardiac | 37 (23%) | 11 (24%) | 26 (22%) | | |
| Unknown | 1 (1%) | 0 (0%) | 1 (1%) | | |
| Baseline PCPC ^c | | | | 0.46 ^b | |
| Normal | 77 (47%) | 23 (51%) | 54 (45%) | | |
| Mild disability | 47 (29%) | 12 (27%) | 35 (29%) | | |
| Moderate disability | 23 (14%) | 7 (16%) | 16 (13%) | | |
| Severe disability | 13 (8%) | 3 (7%) | 10 (8%) | | |
| Coma/vegetative state | 4 (2%) | 0 (0%) | 4 (3%) | | |
| Baseline FSS ^d | 8 [6,11] | 6 [6,9] | 8 [6,12] | 0.02 ^b | |

^a Fisher's exact test.

^b Wilcoxon rank-sum test.

^c Pediatric cerebral performance category.

^d Functional status scale.

^e Day = 7:00AM - 10:59PM. Night = 11:00PM - 6:59AM.

reported as relative risk with 95% confidence intervals. P-values are reported based on a 2-sided alternative and considered statistically significant when less than 0.05. All analyses were performed using SAS software v9.4 (Cary, NC).^{4,12,18–22}

Results

All 164 patients from the original study who met inclusion criteria were included in this secondary analysis. Of the 164 arrests, 45 (27%) occurred at nighttime and 119 (73%) occurred during the daytime. Table 1 describes pre-arrest patient characteristics in the overall cohort, as well as characteristics by time of CPR. Sixty percent of patients were <1 year old; 90 (55%) were male; 82 (50%) were white. Patient pre-existing conditions included 132 (80%) with respiratory insufficiency, 128 (78%) with hypotension, and 99 (60%) with congenital heart disease. Eighty-eight (54%) were surgical cardiac patients (i.e., had cardiac arrest during the post-operative period). Baseline PCPC scores were normal in 77 (47%) patients and 47 (29%) had mild disability. Baseline FSS was lower for patients whose arrest occurred during the nighttime versus those with arrests during daytime (median [IQR]: nighttime: 6 [6,9] vs. daytime: 8 [6,12], p=0.02). Supplemental Table 1 describes pre-arrest patient characteristics by day of week.

Event characteristics for the cohort by time of day and day of week are described in Table 2 and Supplementary Table 2, respectively. The median duration of CPR was 8 min and 100 (61%) events occurred in pediatric cardiac ICUs. The first documented rhythm at the time of CPR was bradycardia with poor perfusion for 91 (55%) of the arrests, followed by 48 (29%) with asystole or PEA. The immediate cause of cardiac arrest was hypotension in 110 (67%) cases. Vasoactive infusions were present at the time of cardiac arrest in 128 (78%) of the cases and there was a non-significant trend toward higher number of vasoactive infusions in place during nighttime arrests (night: 40 (89%) vs. day: 88 (74%), p = 0.06). During CPR, the average DBP was 29.3 mmHg, the average SBP was 74.4, the average chest compression rate was 126 compressions/minute and the average chest compression fraction was 90%. There were no differences in DBP, SBP, chest compression rate, or chest compression fraction between nighttime and daytime arrests.

Table 3 summarizes outcomes by time of day. The DBP thresholds were met during 101/164 (62%) events and SBP thresholds were met during 93/164 (57%) events. Return of spontaneous circulation was achieved in 112/164 (68%) cases and 77/164 (47%) patients survived to hospital discharge. Among patients with ROSC and evaluable post-ROSC BPs, 40/103 (39%) had post-ROSC SBP <5th percentile and nine percent had post-arrest DBP <5th percentile. Post-arrest diastolic BP <5th percentile was nominally more frequent at nighttime (night: 5/37[19%] vs. day: 4/76[5%], p = 0.05). There were no differences in rates

| Table 2 - Event Characteristics by CPR Time of Day. | | | | | | | |
|---|----------------------|-----------------------------|----------------------------|-------------------|--|--|--|
| | Time of CPR | | | | | | |
| | Overall(N = 164) | Night ^c (N = 45) | Day ^c (N = 119) | p-Value | | | |
| Location of CPR Event | | | | 1.00 ^b | | | |
| PICU | 64 (39%) | 18 (40%) | 46 (39%) | | | | |
| CICU | 100 (61%) | 27 (60%) | 73 (61%) | | | | |
| Duration of CPR (min) | 8 [3,27] | 7 [4, 44] | 9 [3,24] | 0.33ª | | | |
| First documented rhythm | | | | 0.90 ^b | | | |
| Asystole/PEA | 48 (29%) | 14 (31%) | 34 (29%) | | | | |
| VF/VT | 19 (12%) | 6 (13%) | 13 (11%) | | | | |
| Bradycardia with pulses | 91 (55%) | 25 (56%) | 66 (55%) | | | | |
| Unknown | 6 (4%) | 0 (0%) | 6 (5%) | | | | |
| Immediate cause | | | | | | | |
| Hypotension | 110 (67%) | 32 (71%) | 78 (66%) | 0.58 ^b | | | |
| Respiratory decompensation | 72 (44%) | 14 (31%) | 58 (49%) | 0.05 ^b | | | |
| Cyanosis without respiratory decompensation | 2 (1%) | 1 (2%) | 1 (1%) | 0.48 ^b | | | |
| Arrhythmia | 31 (19%) | 11 (24%) | 20 (17%) | 0.27 ^b | | | |
| Interventions in place | | | | | | | |
| Central venous catheter | 142 (87%) | 38 (84%) | 104 (87%) | 0.61 ^b | | | |
| Vasoactive infusion | 128 (78%) | 40 (89%) | 88 (74%) | 0.06 ^b | | | |
| Invasive mechanical ventilation | 134 (82%) | 39 (87%) | 95 (80%) | 0.37 ^b | | | |
| Non-invasive ventilation | 19 (12%) | 4 (9%) | 15 (13%) | 0.60 ^b | | | |
| Pharmacologic Interventions | | | | | | | |
| Epinephrine | 143 (87%) | 41 (91%) | 102 (86%) | 0.44 ^b | | | |
| Calcium | 78 (48%) | 21 (47%) | 57 (48%) | 1.00 ^b | | | |
| Sodium bicarbonate | 93 (57%) | 26 (58%) | 67 (56%) | 1.00 ^b | | | |
| Hemodynamics/CPR Mechanics | | | | | | | |
| Average DBP (mmHg) | 29.3 [22.8, 37.9] | 28.3 [25.3, 36.5] | 29.6 [21.8, 38.0] | 0.64 ^a | | | |
| Average SBP (mmHg) | 74.4 [54.9, 98.2] | 77.7 [55.5, 98.2] | 72.7 [54.8, 99.0] | 0.33 ^a | | | |
| Average chest compression rate (/min) | 125.6 [114.3, 139.4] | 120.4 [108.3, 135.5] | 126.7 [115.1, 140.1] | 0.11 ^a | | | |
| Chest compression fraction | 0.9 [0.8, 1.0] | 0.9 [0.9, 1.0] | 0.9 [0.8, 1.0] | 0.69 ^a | | | |
| ^a Wilcovon rank-sum test | | | | | | | |

^b Fisher's exact test.

^c Day = 7:00AM - 10:59PM. Night = 11:00PM - 6:59AM.

| Table 3 - CPR Quality and Outcomes by Time of Day. | | | | | | | |
|--|---------------|--------------------|------------------|-------------------|---|-------------------|--|
| | Time of CPR | | | | Multivariable Regression | | |
| | Overall | Night ^e | Day ^e | P-value | Relative of Risk of Outcome for Night vs. Day (95% C.I.) | p-Value | |
| CPR Quality | | | | | | | |
| DBP Threshold ^a | 101/164 (62%) | 31/45 (69%) | 70/119 (59%) | 0.28 ^h | 1.27 (0.80, 1.98) | 0.30 ⁱ | |
| SBP Threshold ^b | 93/164 (57%) | 25/45 (56%) | 68/119 (57%) | 0.86 ^h | 0.93 (0.56, 1.48) | 0.76 | |
| Compression Rate per Guidelines ^c | 54/164 (33%) | 18/45 (40%) | 36/119 (30%) | 0.26 ^h | 1.19 (0.63, 2.17) | 0.57 ⁱ | |
| Post-arrest hypotension | | | | | | | |
| Any post-arrest systolic BP <5th percentile ^d | 40/103 (39%) | 14/27 (52%) | 26/76 (34%) | 0.12 ^h | 1.54 (0.77, 2.96) | 0.20 ⁱ | |
| Any post-arrest diastolic BP ${<}5\text{th}\ \text{percentile}^{\text{d}}$ | 9/103 (9%) | 5/27 (19%) | 4/76 (5%) | 0.05 ^h | | | |
| Outcomes Immediate Outcome of CPR event | | | | 0.55 ^h | | | |
| $ROSC \ge 20 min$ | 112/164 (68%) | 28/45 (62%) | 84/119 (71%) | | | | |
| ROC with e-CPR | 36/164 (22%) | 12/45 (27%) | 24/119 (20%) | | | | |
| Died | 16/164 (10%) | 5/45 (11%) | 11/119 (9%) | | | | |
| $ROSC \ge 20 min$ | 112/164 (68%) | 28/45 (62%) | 84/119 (71%) | 0.35 ^h | 0.98 (0.61, 1.52) | 0.93 ⁱ | |
| Survival to hospital discharge | 77/164 (47%) | 16/45 (36%) | 61/119 (51%) | 0.08 ^h | 0.72 (0.39, 1.25) | 0.27 ⁱ | |
| 24-h survival | 135/164 (82%) | 35/45 (78%) | 100/119 (84%) | 0.37 ^h | 0.98 (0.64, 1.46) | 0.93 ⁱ | |
| Survival with favorable neurologic outcome | 70/164 (43%) | 14/45 (31%) | 56/119 (47%) | 0.08 ^h | 0.70 (0.37, 1.26) | 0.25 | |
| New morbidity at hospital discharge ^g | 22/77 (29%) | 5/16 (31%) | 17/61 (28%) | 0.77 ^h | 1.09 (0.35, 2.83) | 0.87 ^j | |

 a DBP Threshold defined as mean DBP \geq 25 mmHg for infants; DBP \geq 30 for children \geq 1 year old.

 b SBP Threshold defined as mean SBP \geq 60 mmHg for infants; SBP \geq 80 mmHg for children \geq 1 year old.

^c Compression rate Per Guidelines defined as 100–120 chest compressions/minute.

^d Population: Subjects with post-arrest blood pressure data available. ECMO subjects are excluded.

^e Day = 7:00AM - 10:59PM. Night = 11:00PM - 6:59 AM.

^f Favorable neurologic outcome was defined as no more than moderate disability or no worsening from baseline Pediatric Cerebral Performance Category (PCPC).

⁹ New morbidity is assessed only among survivors to hospital discharge and is defined as an increase in Functional Status Score (FSS) of 3 or greater from baseline to hospital discharge.

h Fisher's exact test.

ⁱ Results are based on Poisson regression models with robust error estimates, adjusting for age category (\leq 1 year, > 1 year), first documented rhythm (asystole/ PEA,VF/VT, bradycardia with poor perfusion), illness category (surgical cardiac, medical cardiac, surgical non-cardiac, medical non-cardiac), location of CPR (PICU, CICU), and study site.

^j Results are based on Poisson regression models with robust error estimates, adjusting for age category (\leq 1 year, > 1 year), first documented rhythm (asystole/PEA,VF/VT, bradycardia with poor perfusion), illness category (surgical cardiac, medical cardiac, surgical non-cardiac, medical non-cardiac), and location of CPR (PICU, CICU).

of ROSC, 24-h survival, survival with favorable neurologic outcome, or new morbidity at hospital discharge between patients who arrested during nighttime versus daytime. There was no difference in survival to hospital discharge among patients who arrested at nighttime (night: 16/ 45 [36%] vs. day: 61/119 [51%], p = 0.08). Supplemental Table 3 summarizes outcomes by day of week.

In Table 4, nighttime and daytime arrests are compared according to whether or not they met intra-arrest DBP targets and whether they had post-arrest systolic or diastolic hypotension. There were no associations between intra-arrest DBP and post-arrest systolic hypotension based on the time of day of cardiac arrest. Similarly, there was no association between intra-arrest DBP and post-arrest diastolic hypotension based on the time of day of cardiac arrest.

Discussion

In this ICU cohort of children with invasive hemodynamic monitoring at the time of in-hospital cardiac arrest, CPR quality was not worse at nighttime compared to daytime, as determined by frequency of attaining DBP thresholds known to be associated with improved outcomes (69% versus 59%). Moreover, there were no differences in median DBPs (28 mmHg [25, 37] vs. 30 mmHg [22, 38]), chest compression rates, or chest compression fractions between groups. In addition, the mean SBP did not differ (78 mmHg [56, 98] versus 73 mmHg [55, 99]), suggesting that the force of chest compressions was similar at nighttime and daytime.²³ Interestingly, diastolic hypotension during the first 20 min post-ROSC was nominally more frequent at nighttime (19% versus 6%, p=0.05), however survival to hospital discharge was not different after nighttime arrests (36% nighttime versus 51% daytime, p=0.08). While previous papers have addressed CPR mechanics at nighttime versus daytime; to our knowledge this is the first study of IHCA that compared CPR hemodynamics at nighttime versus daytime.

Using this prospectively generated database with rich invasive hemodynamic data during CPR, we showed that CPR hemodynamics were not demonstrably different in this highly resourced and monitored cohort of patients that had cardiac arrests in large academic ICUs. The clinical teams were able to provide high-quality CPR regardless of the time of day and achieved DBP ≥ 25 mmHg for

| | Time of CPR | | | | | |
|--|------------------|---------------------------|---------------------------|-------------------|--|--|
| | Overall(N = 103) | Night ^e (N=27) | Day ^e (N = 76) | p-Value | | |
| Intra-arrest DBP Threshold ^b and Post-arrest Systolic Hypotension ^c | | | | 0.23 ^a | | |
| Above DBP threshold and no hypotension | 44/103 (43%) | 9/27 (33%) | 35/76 (46%) | | | |
| Above DBP threshold and hypotension | 26/103 (25%) | 11/27 (41%) | 15/76 (20%) | | | |
| Below DBP threshold and no hypotension | 19/103 (18%) | 4/27 (15%) | 15/76 (20%) | | | |
| Below DBP threshold and hypotension | 14/103 (14%) | 3/27 (11%) | 11/76 (14%) | | | |
| Intra-arrest DBP Threshold ^b and Post-arrest Diastolic Hypotension ^d | | | | 0.08 ^a | | |
| Above DBP threshold and no hypotension | 66/103 (64%) | 17/27 (63%) | 49/76 (64%) | | | |
| Above DBP threshold and hypotension | 4/103 (4%) | 3/27 (11%) | 1/76 (1%) | | | |
| Below DBP threshold and no hypotension | 28/103 (27%) | 5/27 (19%) | 23/76 (30%) | | | |
| Below DBP threshold and hypotension | 5/103 (5%) | 2/27 (7%) | 3/76 (4%) | | | |

Table 4 – Intra-arrest DBP Target and Post-Arrest Hypotension by Time of Day.

^b DBP Threshold defined as mean DBP \geq 25 mmHg for infants; DBP \geq 30 for children \geq 1 year old.

^c Post-arrest Systolic Hypotension defined as average values below the age-based 5th percentile.

^d Post-arrest Diastolic Hypotension defined as average values below the age-based 5th percentile.

^e Day = 7:00AM - 10:59PM. Night = 11:00PM - 6:59 AM.

infants and >30 mmHg for older children in 62% of resuscitations. Esangbedo et al. reported CPR mechanics data in a large cohort of hospitals participating in a CPR quality collaborative (pediRES-Q).⁶ They similarly found no difference in CPR quality indicators including CC depth, CC rate and CCF during the nighttime compared to daytime cardiac arrest. Esangbedo et al. found a significant difference in survival to hospital discharge between the two groups (22% nighttime survival vs. 39% daytime survival) which is consistent with other large registry pediatric studies.⁶ The populations studied were slightly different between the two investigations; with non-ICU arrests included (19% ED and ward) as well as an older population (66% older than 1 year) in the Esangbedo pediRES-Q investigation. The differences in daytime survival between the two studies (51% daytime in the PICqCPR cohort vs. 39% in the pediRES-Q cohort) are possibly accounted for by the younger age, higher percentage of cardiac patients, and inclusion of only ICU arrests in the PICqCPR cohort.^{4,18} Taking both studies into account, two themes emerge: 1) resuscitation parameters (mechanics and hemodynamics of CPR) were not different between nighttime and daytime events and 2) survival to hospital discharge is less likely for arrests that occur during the nighttime.

Why was there no difference in CPR quality at nighttime in either this PICqCPR cohort or the pediRES-Q cohort? Previous investigators postulated that resuscitation practices may be worse at night because of lower staffing ratios and/or provider fatigue at nighttime.^{8,24,25} However, the time of day of cardiac arrest does not impact survival outcomes for subpopulations such as trauma patients or arrests that occur in emergency departments.^{11,26} Peberdy et al. suggested this may be due to consistent staffing of senior residents and attendings for such patients regardless of time of day or day of week.¹¹ While ICU staffing at nighttime may have overall fewer physician team providers, nursing staffing is similar at nighttime and daytime,²⁷ and large academic pediatric ICUs may be able to rally forces around a critical event to briefly provide the same care during CPR as is done during the daytime. Although current overall pediatric physician staffing is lower at night, many hospitals have in-house 24 h pediatric and pediatric cardiac intensivist coverage leading to more senior staffing presence in pediatric ICUs,²⁴ which is apparently sufficient to assure excellent CPR guality.28

The bulk of published data indicate that outcomes from pediatric IHCAs are worse at night, but the reasons for worse outcomes have not been established. Survival to hospital discharge is affected by multiple factors including patient pre-arrest illness, quality of CPR provided, as well as post-cardiac arrest care delivery.18,29 Hypotension, fever, hyperoxia, hypocarbia and severe left ventricular dysfunction in the post-cardiac arrest period are associated with worse neurologic and survival outcomes.^{30–32} Implementation of post-cardiac arrest care is a complex knowledge and time intensive endeavor. Perhaps postcardiac arrest care was less effective at nighttime. Overnight resources may be sufficient to deal with brief acute emergencies, but the apparently less intense and longer duration post-arrest care may be a greater challenge to manage at nighttime. Supporting the contention that the differences in outcomes may be related to differences on postarrest care, post-arrest diastolic hypotension was nominally more frequent at nighttime in this small cohort (19% versus 5%, p=0.051). Because the PICqCPR database did not include post-cardiac arrest care variables beyond the first 20 min post-arrest, we were not able to evaluate the quality of post-arrest care further. Future studies focused on the delivery of post-cardiac arrest care should investigate the quality of post-cardiac arrest care delivered by time of day and its impact on survival to hospital discharge.

These data establish that ICU providers can provide high quality of CPR regardless of the time of day or day of the week. If differences are seen in this quality of CPR at nighttime within a center, quality improvement initiatives may be able to improve the disparity.^{33,34} If high quality CPR is provided, post-arrest care for children with nighttime ICHAs is the next logical focus for potential quality improvement.

This study has several limitations. The relatively small patient cohort available limited the power to demonstrate true differences in survival to hospital discharge at nighttime. The fact that we did not observe survival differences in this study limits our ability to comment on the role of intraarrest quality and hemodynamics in the outcome differences seen between day and night arrests in previous studies. Also, this cohort represents a group of monitored patients in large academic ICUs with arterial lines and may not be generalizable to other ICU patients or patients outside of the ICU setting. However, previous studies have established that the majority of pediatric IHCAs in the United States occur in ICUs, and that arterial lines are often in place at the time of arrest.² For this report we used DBP, SBP, chest compression rate, and chest compression fraction as indicators of CPR quality, but we did not have data regarding chest compression depth, release velocity, or ventilation rate which may be important contributors to outcomes. The median baseline FSS was lower among patients with nighttime versus daytime IHCAs; although this could have conceivably influenced the vigor of resuscitation efforts, there were nevertheless no differences in DBP, SBP, chest compression rate, or chest compression fraction between nighttime and daytime arrests.

Conclusions

In this monitored cohort of pediatric ICU patients who experienced IHCA, CPR quality was not different during nighttime versus daytime. Differences in patient outcomes by time of day are presumably explained by other unmeasured factors to be explored in future research.

Funding support

Supported, in part, by the following cooperative agreements from the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, Department of Health and Human Services: UG1HD050096, UG1HD049981, UG1HD049983, UG1HD063108, UG1HD083171, UG1HD083166, UG1HD083170, U10HD050012, U10HD063106, U10HD063114 and U01HD049934.

Conflicts of interest

The Authors have the following disclosures. Dr. Wolfe has received research funding for NHLBI (funds to institution), as well as speaking honoraria from Zoll Medical. Robert A Berg received research grants from NHLBI and NICHD to study pediatric in-hospital cardiopulmonary resuscitation (funds to institution), as well as speaking honoraria from Japan Pediatric Society for lecture on pediatric cardiopulmonary resuscitation. Dr. Sutton has received research grants from NHLBI and NICHD to study pediatric in-hospital cardiopulmonary resuscitation (funds to institution), Pediatric Advance Life Support authorship via the American Heart Association, membership on the Emergency Cardiovascular Care Committee and past speaking honoraria for Zoll Medical. The rest of the authors report no COI.

Appendix A. CPCCRN PICqCPR Investigators

Athena F. Zuppa^a, Katherine Graham^a, Carolann Twelves^a, Mary Ann Diliberto^a, Richard Holubkov^b, Russell Telford^b, Christopher Locandro^b, Whitney Coleman^b, Alecia Peterson^b, Julie Thelen^b, Sabrina Heidemann^c, Ann Pawluszka^c, Elyse Tomanio^d, Michael J. Bell^{d,g}, Mark W. Hall^e, Lisa Steele^e, Jeni Kwok^f, Anil Sapru^{f,h}, Alan Abraham^g, Mustafa F. Alkhouli^h, Thomas P. Shanleyⁿ, Monica Weber^j, Heidi J. Dalton^k, Aimee La Bell^I, Peter M. Mourani^I, Kathryn Malone^I, Allan Doctor^o

^aDepartment of Anesthesiology and Critical Care Medicine, The Children's Hospital of Philadelphia, University of Pennsylvania, Philadelphia, PA, United States ^bDepartment of Pediatrics, University of Utah, Salt Lake City, Utah, United States

^cDepartment of Pediatrics, Children's Hospital of Michigan, Wayne State University, Detroit, MI, United States

^dDepartment of Pediatrics, Children's National Medical Center, Washington D.C., United States

^eDepartment of Pediatrics, Nationwide Children's Hospital, The Ohio State University, Columbus, OH, United States

^fDepartment of Anesthesiology, Children's Hospital Los Angeles, University of Southern California Keck School of Medicine, Los Angeles, CA, United States

^gDepartment of Critical Care Medicine, Children's Hospital of Pittsburgh, University of Pittsburgh, Pittsburgh, PA, United States

^hDepartment of Pediatrics, Benioff Children's Hospital, University of California San Francisco, San Francisco, CA, United States

ⁱDepartment of Pediatrics, Mattel Children's Hospital, University of California Los Angeles, Los Angeles, CA, United States

^jDepartment of Pediatrics, C.S. Mott Children's Hospital, University of Michigan, MI, United States

^kDepartment of Pediatrics, Phoenix Children's Hospital, Phoenix, AZ, United States

^IDepartment of Pediatrics, Denver Children's Hospital, University of Colorado, Denver, CO, United States

^mDepartment of Molecular Biology, Princeton University, Princeton, New Jersey, United States

ⁿDepartment of Pediatrics, Lurie Children's Hospital, Northwestern University, Chicago, IL, United States

^oDepartment of Pediatrics, University of Maryland, Baltimore, MD, United States

Appendix B. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:https://doi.org/10.1016/j.resuscitation.2020.06.027.

REFERENCES

- Holmberg MJ, Ross CE, Fitzmaurice GM, et al. Annual incidence of adult and pediatric in-hospital cardiac arrest in the United States. Circ Cardiovasc Qual Outcomes 2019;12:1–8, doi:http://dx.doi.org/ 10.1161/circoutcomes.119.005580.
- Berg RA, Sutton RM, Holubkov R, et al. Ratio of PICU versus ward cardiopulmonary resuscitation events is increasing. Crit Care Med 2013;41(October (10)):2292–7, doi:http://dx.doi.org/10.1097/ CCM.0b013e31828cf0c0.
- Gupta P, Tang X, Gall CM, Lauer C, Rice TB, Wetzel RC. Epidemiology and outcomes of in-hospital cardiac arrest in critically ill children across hospitals of varied center volume: a multi-center analysis. Resuscitation 2014;85:1473–9, doi:http://dx.doi.org/ 10.1016/j.resuscitation.2014.07.016.
- Lowry AW, Knudson JD, Cabrera AG, Graves DE, Morales DLS, Rossano JW. Cardiopulmonary resuscitation in hospitalized children with cardiovascular disease: estimated prevalence and outcomes from the Kids' Inpatient database. Pediatr Crit Care Med 2013;14:248 –55, doi:http://dx.doi.org/10.1097/PCC.0b013e3182713329.
- Girotra S, Spertus JA, Li Y, Berg RA, Nadkarni VM, Chan PS. Survival trends in pediatric in-hospital cardiac arrests an analysis from get with the guidelines-resuscitation. Circ Cardiovasc Qual Outcomes 2013;6:42–9, doi:http://dx.doi.org/10.1161/ CIRCOUTCOMES.112.967968.

- Esangbedo I, Yu P, Raymond T, et al. Pediatric in-hospital CPR quality at night and on weekends. Resuscitation 2020;146:56–63, doi:http:// dx.doi.org/10.1016/j.resuscitation.2019.10.039.
- Karlsson LIM, Wissenberg M, Fosbøl EL, et al. Diurnal variations in incidence and outcome of out-of-hospital cardiac arrest including prior comorbidity and pharmacotherapy: a nationwide study in Denmark. Resuscitation 2014;85(September (9)):1161–8, doi:http://dx.doi.org/ 10.1016/j.resuscitation.2014.06.012.
- Bhanji F, Topjian AA, Nadkarni VM, et al. Survival rates following pediatric in-hospital cardiac arrests during nights and weekends. JAMA Pediatr 2017;171:39–45, doi:http://dx.doi.org/10.1001/ jamapediatrics.2016.2535.
- Kitamura T, Kiyohara K, Nitta M, Nadkarni VM, Berg RA, Iwami T. Survival following witnessed pediatric out-of-hospital cardiac arrests during nights and weekends. Resuscitation 2014;85(December (12)):1692–8, doi:http://dx.doi.org/10.1016/j. resuscitation.2014.08.035.
- Bagai A, McNally BF, Al-Khatib SM, et al. Temporal differences in outof-hospital cardiac arrest incidence and survival. Circulation 2013;128:2595–602, doi:http://dx.doi.org/10.1161/ CIRCULATIONAHA.113.004164.
- Peberdy MA, Ornato JP, Larkin GL, et al. Survival from in-hospital cardiac arrest during nights and weekends. JAMA - J Am Med Assoc 2008;299:785–92, doi:http://dx.doi.org/10.1001/jama.299.7.785.
- 12. Jayaram N, Spertus JA, Nadkarni V, et al. Hospital variation in survival after pediatric in-hospital cardiac arrest. Circ Cardiovasc Qual Outcomes 2014;7:517–23, doi:http://dx.doi.org/10.1161/CIRCOUTCOMES.113.000691.
- Berg RA, Sutton RM, Reeder RW, et al. Association between diastolic blood pressure during pediatric in-hospital cardiopulmonary resuscitation and survival. Circulation 2017137:, doi:http://dx.doi.org/ 10.1161/CIRCULATIONAHA.117.032270 CIRCULATIONAHA.117.032270.
- 14. Jacobs I, Nadkarni V, Bahr J, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries. A statement for healthcare professionals from a task force of the International Liaison Committee on Resusci. Circulation 2004;110:3385–97, doi:http://dx.doi.org/ 10.1161/01.CIR.0000147236.85306.15.
- 15. Nolan JP, Berg RA, Andersen LW, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein Resuscitation Registry template for in-hospital cardiac arrest: a Consensus Report From a Task Force of the International Liaison Committee on Resuscitation (American). Circulation 2019;140:e746 -57, doi:http://dx.doi.org/10.1161/CIR.000000000000710.
- Pollack MM, Holubkov R, Glass P, et al. Functional status scale: new pediatric outcome measure. Pediatrics 2009;124(July (1)):e18–28, doi:http://dx.doi.org/10.1542/peds.2008-1987.
- Rosner B, Cook N, Portman R, Daniels S, Falkner B. Determination of blood pressure percentiles in normal-weight children: some methodological issues. Am J Epidemiol 2008;167(March (6)):653–66, doi:http://dx.doi.org/10.1093/aje/kwm348.
- Meaney Pa, Nadkarni VM, Cook EF, et al. Higher survival rates among younger patients after pediatric intensive care unit cardiac arrests. Pediatrics 2006;118:2424–33, doi:http://dx.doi.org/10.1542/ peds.2006-1724.
- Nadkarni VM, Larkin GL, Peberdy MA, et al. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. J Am Med Assoc 2006;295:50–7, doi:http://dx.doi.org/ 10.1001/jama.295.1.50.

- Greenland S. Model-based estimation of relative risks and other epidemiologic measures in studies of common outcomes and in casecontrol studies. Am J Epidemiol 2004;160:301–5, doi:http://dx.doi. org/10.1093/aje/kwh221.
- Gupta P, Rettiganti M, Jeffries HE, et al. Risk factors and outcomes of in-hospital cardiac arrest following pediatric heart operations of varying complexity. Resuscitation 2016;105:1–7, doi:http://dx.doi. org/10.1016/j.resuscitation.2016.04.022.
- 22. Matos RI, Watson RS, Nadkarni VM, et al. Duration of cardiopulmonary resuscitation and illness category impact survival and neurologic outcomes for in-hospital pediatric cardiac arrests. Circulation 2013;127:442–51, doi:http://dx.doi.org/10.1161/ CIRCULATIONAHA.112.125625.
- Sainio M, Sutton RM, Huhtala H, et al. Association of arterial blood pressure and CPR quality in a child using three different compression techniques, a case report. Scand J Trauma Resusc Emerg Med 2013;21:1–6, doi:http://dx.doi.org/10.1186/1757-7241-21-51.
- 24. Gupta P, Rettiganti M, Rice TB, Wetzel RC. Impact of 24/7 in-hospital intensivist coverage on outcomes in pediatric intensive care a multicenter study. Am J Respir Crit Care Med 2016;194(12):1506–13, doi:http://dx.doi.org/10.1164/rccm.201512-24560C.
- McHugh MD, Rochman MF, Sloane DM, et al. Better nurse staffing and nurse work environments associated with increased survival of inhospital cardiac arrest patients. Med Care 2016;54(January (1)): 74–80, doi:http://dx.doi.org/10.1097/MLR.000000000000456.
- 26. Carr BG, Jenkins P, Branas CC, et al. Does the trauma system protect against the weekend effect? J Trauma 2010;69(November (5)): 1042–7, doi:http://dx.doi.org/10.1097/TA.0b013e3181f6f958.
- Needleman J, Buerhaus P, Mattke S, Stewart M, Zelevinsky K. Nursestaffing levels and the quality of care in hospitals. N Engl J Med 2002;346:1715–22, doi:http://dx.doi.org/10.1056/NEJMsa012247.
- Wheeler DS, Dewan M, Maxwell A, Riley CL, Stalets EL. Staffing and workforce issues in the pediatric intensive care unit. Transl Pediatr 2018;7:275–83, doi:http://dx.doi.org/10.21037/tp.2018.09.05.
- Sutton RM, French B, Niles DE, et al. 2010 American Heart Association recommended compression depths during pediatric inhospital resuscitations are associated with survival. Resuscitation 2014;85:1179–84, doi:http://dx.doi.org/10.1016/j. resuscitation.2014.05.007.
- Topjian AA, French B, Sutton RM, et al. Early postresuscitation hypotension is associated with increased mortality following pediatric cardiac arrest. Crit Care Med 2014;42:1518–23, doi:http://dx.doi.org/ 10.1097/CCM.00000000000216.
- Conlon TW, Falkensammer CB, Hammond RS, Nadkarni VM, Berg RA, Topjian AA. Association of left ventricular systolic function and vasopressor support with survival following pediatric out-of-hospital cardiac arrest. Pediatr Crit Care Med 2015;16(February (2)):146–54, doi:http://dx.doi.org/10.1097/PCC.00000000000305.
- 32. Gebhardt K, Guyette FX, Doshi AA, et al. Prevalence and effect of fever on outcome following resuscitation from cardiac arrest. Resuscitation 2013;84:1062–7, doi:http://dx.doi.org/10.1016/j. resuscitation.2013.03.038.
- Wolfe H, Zebuhr C, Topjian AA, et al. Interdisciplinary ICU cardiac arrest debriefing improves survival outcomes. Crit Care Med 2014;42:1688–95, doi:http://dx.doi.org/10.1097/ CCM.00000000000327.
- 34. Hunt EA, Jeffers J, McNamara L, et al. Improved cardiopulmonary resuscitation performance with CODE ACES²: A resuscitation quality bundle. J Am Heart Assoc 20187:, doi:http://dx.doi.org/10.1161/ JAHA.118.009860.